

*陳瓊珠醫師

所有發表期刊論文

1. *Chen CC, Lin WY, Chan HL, Tu PH, Lee ST, Lu CS. Brown P The impact of low-frequency stimulation of subthalamic region on self-generated isometric contraction in patients with Parkinson's disease. *Exp Brain Res.* 2013;227:53-62.
2. Wu-Chou YH, Chen YT, Yeh TH, Chang HC, Weng YH, Lai SC, Huang CL, Chen RS, Huang YZ, *Chen CC, Hung J, Chuang WL, Lin WY, Chen CH, Lu CS. Genetic variants of SNCA and LRRK2 genes are associated with sporadic PD susceptibility: a replication study in a Taiwanese cohort. *Parkinsonism Relat Disord.* 2013;19:251-255.
3. Hsu YT, Lai HY, Chang YC, Chiou SM, Lu MK, Lin YC, Liu YL, *Chen CC, Huang HC, Chien TF, Lin SZ, Chen YY, Tsai CH. The role of the sub-thalamic nucleus in the preparation of volitional movement termination in Parkinson's disease. *Exp Neurol.* 2012;233:253-263.
4. *Chen CC, Lin WY, Chan HL, Hsu YT, Tu PH, Lee ST, Chiou SM, Tsai CH, Lu CS and Brown P. Stimulation of the subthalamic region at 20 Hz slows the development of grip force in Parkinson's disease. *Exp Neurol.* 2011;231:91-96.
5. Chen P, Lin JJ, Lu CS, Ong CT, Hsieh PF, Yang CC, Tai CT, Wu SL, Lu CH, Hsu YC, Yu HY, Ro LS, Lu CT, *Chu CC, Tsai JJ, Su YH, Lan SH, Sung SF, Lin SY, Chuang HP, Huang LC, Chen YJ, Tsai PJ, Liao HT, Lin YH, Chen CH, Chung WH, Hung SI, Wu JY, Chang CF, Chen L, Chen YT, Shen CY Carbamazepine-induced toxic effects and HLA-B*1502 screening in Taiwan. *N Engl J Med.* 2011;364:1126-1133.
6. *Chen CC, Hsu YT, Chan HL, Chiou SM, Tu PH, Lee ST, Tsai CH, Lu CS. Brown P Complexity of subthalamic 13-35 Hz oscillatory activity directly correlates with clinical impairment in patients with Parkinson's disease. *Exp Neurol.* 2010;224:234-240.
7. Pogosyan A, Yoshida F, *Chen CC, Martinez-Torres I, Foltynie T, Limousin P, Zrinzo L, Hariz MI, Brown P. Parkinsonian Impairment Correlates With Spatially Extensive Subthalamic Oscillatory Synchronisation. *Neuroscience.* 2010;171:245-257.

8. Lai SC, Jung SM, Grattan-Smith P, Sugo E, Lin YW, Chen RS, *Chen CC, Wu-Chou YH, Lang AE, Lu CS Neuronal intranuclear inclusion disease: two cases of dopa-responsive juvenile parkinsonism with drug-induced dyskinesia. *Mov Disord.* 2010;25:1274-1279.
9. Yoshida F, Martinez-Torres I, Pogosyan A, Holl E, Petersen E, *Chen CC, Foltynie T, Limousin P, Zrinzo LU, Hariz MI, Brown P. Value of subthalamic nucleus local field potentials recordings in predicting stimulation parameters for deep brain stimulation in Parkinson's disease. *J Neurol Neurosurg Psychiatry.* 2010;81:885-889.
10. Eusebio A, *Chen CC, Lu CS, Lee ST, Tsai CH, Limousin P, Hariz M, Brown P. Effects of low-frequency stimulation of the subthalamic nucleus on movement in Parkinson's disease. *Exp Neurol.* 2008;209:125-130.
11. *Chen CC, Brown P. The role of synchronized low frequency activity in globus pallidus interna in dystonia. *Acta Neurol Taiwan.* 2007;16:1-6.
12. *Chen CC, Litvak V, Gilbertson T, Kuhn AA, Lu CS, Lee ST, Tsai CH, Tisch S, Limousin P, Hariz M, Brown P. Excessive synchronization of basal ganglia neurons at 20 Hz slows movement in Parkinson's disease. *Exp Neurol.* 2007;205:214-221.
13. Brucke C, Kupsch A, Schneider GH, Hariz MI, Nuttin B, Kopp U, Kempf F, Trottenberg T, Doyle L, *Chen CC, Yarrow K, Brown P, Kuhn AA. The subthalamic region is activated during valence-related emotional processing in patients with Parkinson's disease. *Eur J Neurosci.* 2007;26:767-774.
14. Androulidakis AG, Kuhn AA, *Chen CC, Blomstedt P, Kempf F, Kupsch A, Schneider GH, Doyle L, Dowsey-Liousin P, Hariz MI, Brown P. Dopaminergic therapy promotes lateralized motor activity in the subthalamic area in Parkinson's disease. *Brain* 2007;130:457-468.
15. Kempf F, Brucke C, Kuhn AA, Schneider GH, Kupsch A, *Chen CC, Androulidakis AG, Wang S, Vandenberghe W, Nuttin B, Aziz T, Brown P. Modulation by dopamine of human basal ganglia involvement in feedback control of movement. *Curr Biol.* 2007;17:587-589.
16. *Chen CC, Pogosyan A, Zrinzo LU, Tisch S, Limousin P, Ashkan K, Yousry T,

Hariz MI, Brown P. Intra-operative recordings of local field potentials can help localize the subthalamic nucleus in Parkinson's disease surgery. *Exp Neurol*. 2006;198:214-221.

17. *Chen CC, Kuhn AA, Hoffmann KT, Kupsch A, Schneider GH, Trottenberg T, Krauss JK, Wohrle JC, Bardinet E, Yelnik J, Brown P. Oscillatory pallidal local field potential activity correlates with involuntary EMG in dystonia. *Neurology*. 2006;66:418-420.
18. *Chen CC, Brucke C, Kempf F, Kupsch A, Lu CS, Lee ST, Tisch S, Limousin P, Hariz M, Brown P. Deep brain stimulation of the subthalamic nucleus: a two edged sword. *Curr Biol* 2006;16:952-953.
19. *Chen CC, Kuhn AA, Trottenberg T, Kupsch A, Schneider GH, Brown P. Neuronal activity in globus pallidus interna can be synchronized to local field potential activity over 3-12 Hz in patients with dystonia. *Exp Neurol*. 2006;202:480-486.
20. Brown P, *Chen CC, Wang S, Kuhn AA, Doyle L, Yarrow K, Nuttin B, Stein J, Aziz T. Involvement of human basal ganglia in off-line feed-back control of voluntary movement. *Curr Biol*. 2006;16:2129-2134.
21. Momeni P, Lu CS, Chou YH, Chang HC, Chen RS, *Chen CC, Hsu JT, Singleton A, Hardy J. Taiwanese cases of SCA2 are derived from a single founder. *Movement Disorders* 2005;20:1633-1636.
21. Yen TH, Lai PC, *Chen CC, Hsueh S, Huang JY. Renal involvement in patients with polymyositis and dermatomyositis. *Int J Clin Pract*. 2005;59:188-193.
23. *Chen CC, Lee ST, Wu T, Chen CJ, Chen MC, Lu CS. Short-term effect of bilateral subthalamic stimulation for advanced Parkinson's disease. *Chang Gung Med J*. 2003;26:344-351.
24. *Chen CC, Lee ST, Wu T, Chen CJ, Huang CC, Lu CS. Hemiballism after subthalamotomy in patients with Parkinson's disease: report of 2 cases. *Mov Disord*. 2002;17:1367-1371.s