

*李佩賢醫師

所有發表期刊論文

1. [Cannabinoid receptor 1 disturbance of PPAR \$\gamma\$ 2 augments hyperglycemia induction of mesangial inflammation and fibrosis in renal glomeruli.](#)

Lin CL, Hsu YC, Lee PH, Lei CC, Wang JY, Huang YT, Wang SY, Wang FS.

J Mol Med (Berl). 2014 Jul;92(7):779-92. doi: 10.1007/s00109-014-1125-6. Epub 2014 Apr 11.

2. [MicroRNA-29a promotion of nephrin acetylation ameliorates hyperglycemia-induced podocyte dysfunction.](#)

Lin CL, Lee PH, Hsu YC, Lei CC, Ko JY, Chuang PC, Huang YT, Wang SY, Wu SL, Chen YS, Chiang WC, Reiser J, Wang FS.

J Am Soc Nephrol. 2014 Aug;25(8):1698-709. doi: 10.1681/ASN.2013050527. Epub 2014 Feb 27.

3. [Analgesic use, parents' clan, and coffee intake are three independent risk factors of chronic kidney disease in middle and elderly-aged population: a community-based study.](#)

Hsu YC, Lee PH, Lei CC, Shih YH, Lin CL.

Ren Fail. 2014 Apr;36(3):361-6. doi: 10.3109/0886022X.2013.866017. Epub 2014 Feb 10.

4. [Hyperuricemia is an independent risk factor for new onset micro-albuminuria in a middle-aged and elderly population: a prospective cohort study in taiwan.](#)

Chang HY, Lee PH, Lei CC, Tung CW, Hsu YC, Huang TJ, Lu LC, Lin CL.

PLoS One. 2013 Apr 24;8(4):e61450. doi: 10.1371/journal.pone.0061450. Print 2013.

5. [Educational intervention in CKD retards disease progression and reduces medical costs for patients with stage 5 CKD.](#)

Lei CC, Lee PH, Hsu YC, Chang HY, Tung CW, Shih YH, Lin CL.

Ren Fail. 2013;35(1):9-16. doi: 10.3109/0886022X.2012.731997. Epub 2012 Oct 22.

6. [Sustained Wnt/ \$\beta\$ -catenin signaling rescues high glucose induction of transforming growth factor- \$\beta\$ 1-mediated renal fibrosis.](#)

Ho C, Lee PH, Hsu YC, Wang FS, Huang YT, Lin CL.

Am J Med Sci. 2012 Nov;344(5):374-82. doi: 10.1097/MAJ.0b013e31824369c5.

7. [Obstructive sleep apnea: a stand-alone risk factor for chronic kidney disease.](#)

Chou YT, Lee PH, Yang CT, Lin CL, Veasey S, Chuang LP, Lin SW, Lin YS, Chen NH.

Nephrol Dial Transplant. 2011 Jul;26(7):2244-50. doi: 10.1093/ndt/gfq821. Epub 2011 Feb 11.

8. [Hyperuricemia as an independent risk factor of chronic kidney disease in middle-aged and elderly population.](#)

Chang HY, Tung CW, Lee PH, Lei CC, Hsu YC, Chang HH, Yang HF, Lu LC, Jong MC, Chen CY, Fang KY, Chao YS, Shih YH, Lin CL.

Am J Med Sci. 2010 Jun;339(6):509-15. doi: 10.1097/MAJ.0b013e3181db6e16.

9. [Hypertriglyceridemia: an independent risk factor of chronic kidney disease in Taiwanese adults.](#)

Lee PH, Chang HY, Tung CW, Hsu YC, Lei CC, Chang HH, Yang HF, Lu LC, Jong MC, Chen CY, Fang KY, Chao YS, Shih YH, Lin CL.

Am J Med Sci. 2009 Sep;338(3):185-9. doi: 10.1097/MAJ.0b013e3181a92804.

10. [Ras activation modulates methylglyoxal-induced mesangial cell apoptosis through superoxide production.](#)

Huang WJ, Tung CW, Ho C, Yang JT, Chen ML, Chang PJ, Lee PH, Lin CL, Wang JY.

Ren Fail. 2007;29(7):911-21.

11. [Methylglyoxal-induced fibronectin gene expression through Ras-mediated NADPH oxidase activation in renal mesangial cells.](#)

Ho C, Lee PH, Huang WJ, Hsu YC, Lin CL, Wang JY.

Nephrology (Carlton). 2007 Aug;12(4):348-56.

12. [Octreotide therapy for chylous ascites in a chronic dialysis patient.](#)

Lee PH, Lin CL, Lai PC, Yang CW.

Nephrology (Carlton). 2005 Aug;10(4):344-7.